Special Site Dermoscopy: Volar Skin (palmar & plantar surfaces)

Ashfaq A. Marghoob, M.D. Attending Physician










Disclosures


- American Dermoscopy Meeting
- Canfield
- Casio
- DermLite
- FotoFinder
- Heine
- HealthCert
- Lecturing
- Publications: Books
- UpToDate




Microsoft Bing AI image generator (no need for copyright)

Dermatoglyphics


Micro-anatomy







T. Saida et al. Clin.Dermatol. 20 (3):279-285, 2002.

Ridges & Furrows




Anatomy







Why is it so important to locate ridges and furrows?


Anatomy of Volar skin



Parallel furrow pattern










- Secretory portion of eccrine sweat glands provide an anatomical niche for melanocyte-melanoma precursor cells
- This explains preferential distribution of early melanoma cells around sweat glands (crista profunda intermedia) in human volar skin = parallel ridge pattern


Okamoto et al. 2014


Parallel ridge pattern

















Both are Melanoma specific patterns

Significance of Dermoscopic Patterns in Detecting Malignant Melanoma on Acral Volar Skin

Results of a Multicenter Study in Japan

Toshiaki Saida, MD, PhD; Atsushi Miyazaki, MD; Shinji Oguchi, MD, PhD; Yasushi Ishihara, MD; Yoriko Yamazaki, MD; Sumio Murase, MD, PhD; Shusuke Yoshikawa, MD; Tetsuya Tsuchida, MD, PhD; Yasuhiro Kawabata, MD, PhD; Kunihiko Tamaki, MD, PhD

Objective: To determine diagnostic variables such as sensitivity and specificity of the major dermoscopic patterns observed in melanocytic lesions on acral volar skin, with particular attention to the significance of the parallel ridge pattern and irregular diffuse pigmentation in detecting acral melanoma.

Design: Multicenter, retrospective study.

Setting: University hospitals in Japan.

Patients: Patients with melanocytic lesions on acral volar skin. A total of 712 melanocytic lesions (103 malignant melanomas, including 36 in situ lesions, and 609 melanocytic nevi) were consecutively collected from the files of 3 hospitals. Diagnoses of all the lesions had been determined histopathologically.

Interventions: Dermoscopic examination.


Main Outcome Measures: The sensitivity, specificity, positive predictive value, negative predictive value,

and diagnostic accuracy of the major dermoscopic patterns seen in benign and malignant melanocytic lesions on acral volar skin.

Results: The parallel ridge pattern and irregular diffuse pigmentation showed extremely high specificity (99.0% and 96.6%, respectively) and very high negative predictive value (97.7% and 97.5%, respectively) in malignant melanoma. For melanoma in situ, the positive predictive value and diagnostic accuracy of the parallel ridge pattern were significantly higher than those of irregular diffuse pigmentation (P=.009 and P=.006, respectively). In melanocytic nevi, the specificity and positive predictive value of the parallel furrow pattern and/or the latticelike pattern were found to be very high (93.2% and 98.3%, respectively).


Conclusions: Dermoscopy is immensely helpful in differentiating malignant melanomas from melanocytic nevi on acral volar skin. Moreover, the parallel ridge pattern aids in detecting acral melanomas in early, curable stages.


Arch Dermatol. 2004;140:1233-1238







Parallel ridge pattern 98% of melanomas <1% volar nevi















Exceptions to the PRP rule for MM

- 1. Peutz-Jeghers syndrome macules
- 2. Laugier-Hunziker syndrome
- 3. Ethnic pigmented macules: progressive volar melanotic macules (PVMM)
- 4. Dye (exogenous)
- 5. Chemotherapy induced pigmentation
- 6. CMN (<1% of other nevi)
- 7. Subcorneal hemorrhage

Benign Dermoscopic Parallel Ridge Pattern Variants

Alice Phan, MD; Stéphane Dalle, MD; Marie-Cécile Marcilly, MD; Jean-Pierre Bergues, MD; Luc Thomas, MD, PhD; Centre Hospitalier Lyon-Sud, Claude Bernard University, Pierre Bénite, France


3. Ethnic pigmented macules: progressive volar melanotic macules (PVMM)





My experience: Context is key. These darker macules occur in a background of normal skin that also has a PRP!















Irregular Diffuse Pigmentation Multiple shades of brown, blue/black




Irregular (asymmetric, disorganized, or multi-colored) diffuse pigmentation (blotch) that encompasses fissures & ridges










Exceptions to IDP rule for MM


- 1. Blue nevus
 - homogeneous blue crossing ridges & furrows
 - can display multiple shades of blue & brown
- 2. Congenital nevus (tardive)
 - parallel furrow pattern with dots/globules on ridges
 - By definition they display a multicomponent pattern


- Blue nevi on volar surfaces cross ridges & furrows
- Often display slight IDP (histo vs biology)



- CMN tend to display a unique multicomponent pattern know as "peas-in-the-pod" pattern
- will discuss together with the other benign patterns..







Validation of a dermatoscopy-based algorithm for the diagnosis of acral melanoma

Müller C, Kittler H, Tschandl P, Rinner C, Grausenburger M-L, Kyrgidis A, Koga H, Moscarella E, Apalla Z, Di Stefani A, Kobayashi K, Lazaridou E, Longo C, Phan A, Saida T, Sotiriou E, Tanaka M, Thomas L, Zalaudek I, Argenziano G, Lallas A

Simplified checklist

parallel ridge pattern


irregular blotch


asymmetry of colors


asymmetry of structures


Sensitivity 95% Specificity 60%


IDP


















Parallel-furrow pattern

linear pigmentation in the furrows



Dermoscopy patterns	Definition	Schema
Benign patterns		
Parrallel furrow pattern	Pigmentation following the furrows	- • • •





9% of melanomas (focal with PRP/IDP)



42% of nevi (throughout & predominant pattern)











• Parallel furrow pattern

a) single line variant sulcus → ridge [


c) double line variant



b) single dotted line variant



b) double dotted line variant










Dermoscopic Characteristics of Congenital Melanocytic Nevi Affecting Acral Volar Skin


Akane Minagawa, MD; Hiroshi Koga, MD; Toshiaki Saida, MD, PhD









Anatomy




Nevomelanocytic nests surrounding adnexal structures






Anatomy



Nevomelanocytic nests in the papillary dermis

Anatomy



Nevomelanocytic nests in the papillary dermis

Dermoscopic Characteristics of Congenital Melanocytic Nevi Affecting Acral Volar Skin

Akane Minagawa, MD; Hiroshi Koga, MD; Toshiaki Saida, MD, PhD

Tardive CMN

2004 through February 28, 2000, we selected cases with

pic features of

uation of der-

rsity hospital.

om January 1, t were present months of life.


ristic dermo-


CMN lesions. a combination lel furrow patar pattern. The owing single-

component patterns: the parallel furrow pattern in 6 (25%), the crista dotted pattern in 3 (12%), the fibrillar pattern in 2 (8%), and the globular, globulostreaklike, nontypical, and parallel ridge patterns in 1 each (4%). We also followed up 6 lesions for several years. Changes in the dermoscopic features were observed in 4 CMN lesions from patients younger than 14 years. Three lesions had a combination of the crista dotted and parallel furrow patterns on the first visit that changed to the nontypical pattern; in addition, the degrees of pigmentation decreased during follow-up. In the remaining lesion, the globulostreaklike pattern changed to the parallel furrow pattern.

Conclusions: Most CMN lesions affecting acral volar skin show characteristic dermoscopic features distinguishable from acral melanoma. The combination of the crista dotted and parallel furrow patterns (ie, peas-in-a-pod pattern) is the most common feature in acral CMN. Some lesions of acral CMN fade during childhood.


Arch Dermatol. 2011;147(7):809-813






Lattice-like pattern


- linear pigmentation in the furrows
- with cross striations across the ridges





Lattice-like pattern











Significance of Dermoscopic Patterns in Detecting Malignant Melanoma on Acral Volar Skin

Results of a Multicenter Study in Japan

Toshiaki Saida, MD, PhD; Atsushi Miyazaki, MD; Shinji Oguchi, MD, PhD; Yasushi Ishihara, MD; Yoriko Yamazaki, MD; Sumio Murase, MD, PhD; Shusuke Yoshikawa, MD; Tetsuya Tsuchida, MD, PhD; Yasuhiro Kawabata, MD, PhD; Kunihiko Tamaki, MD, PhD

Objective: To determine diagnostic variables such as sensitivity and specificity of the major dermoscopic patterns observed in melanocytic lesions on acral volar skin, with particular attention to the significance of the parallel ridge pattern and irregular diffuse pigmentation in detecting acral melanoma.

Design: Multicenter, retrospective study.

Setting: University hospitals in Japan.

Patients: Patients with melanocytic lesions on acral volar skin. A total of 712 melanocytic lesions (103 malignant melanomas, including 36 in situ lesions, and 60 melanocytic nevi) were consecutively collected from the files of 3 hospitals. Diagnoses of all the lesions had been determined histopathologically.

Interventions: Dermoscopic examination.


Main Outcome Measures: The sensitivity, specificity, positive predictive value, negative predictive value,

and diagnostic accuracy of the major dermoscopic patterns seen in benign and malignant melanocytic lesions on acral volar skin.

Results: The parallel ridge pattern and irregular diffuse pigmentation showed extremely high specificity (99.0% and 96.6%, respectively) and very high negative predictive value (97.7% and 97.5%, respectively) in malignant melanoma. For melanoma in situ, the positive predictive value and diagnostic accuracy of the parallel ridge pattern were significantly higher than those of irregular diffuse pigmentation (P=.009 and P=.006, respectively). In melanocytic nevi, the specificity and positive predictive value of the parallel furrow pattern and/or the latticelike pattern were found to be very high (93.2% and 98.3%, respectively).


Conclusions: Dermoscopy is immensely helpful in differentiating malignant melanomas from melanocytic nevi on acral volar skin. Moreover, the parallel ridge pattern aids in detecting acral melanomas in early, curable stages.

Arch Dermatol. 2004;140:1233-1238



Fibrillar pattern


Lines crossing at an angle across both the furrows and ridges





Histology fibrillar pattern:

Weight bearing sole




column slants


Ref: Myazaki A et al.: JAAD 53:230, 2005


Fibrillar pattern: predominant pattern throughout

35% of melanomas 33% of nevi











Benign Dermoscopic volar patterns

Pattern	Soles N=165	Palms N=45	Total N=210
"Lattice-like"	17 (10%)	9 (20%)	26 (12%)
"Fibrillar"	13 (7,9%)	0	13 (6%)

J. Malvehy, S. Puig. Dermoscopic patterns of benign volar melanocytic lesions in patients with atypical mole syndrome. Arch Dermatol, 2004








Fibrillar pattern on plantar surface: Typical vs Atypical


 The key is to determine if the fibrillar pattern is anchored in furrows or on the ridges

 The morphology of the fibrillar pattern can also be helpful (organized vs disorganized)



Typical Fibrillar pattern: anchored in furrows & organized

Typical Atypical



Junctional/compound nevus



Melanoma in situ




Dermatology 2009;218:385-386


DOI: 10.1159/000202986

Oblique View Dermoscopy Changes Regular Fibrillar Pattern into Parallel Furrow Pattern


Yoshifumi Maumi, Masayuki Kimoto, Ken Kobayashi, Naoko Ito, Toshiaki Saida, Masaru Tanaka

Department of Dermatology, Tokyo Women's Medical University Medical Center East, Tokyo, and Shinshu University School of Medicine, Matsumoto, Japan





Fibrillar pattern anchored in the furrows (Type A)




"Horizontal pressure"



At rest: fibrillar


Horizontal pressure: furrow












Atypical Fibrillar pattern: anchored in ridges & disorganized


Typical



Junctional/compound nevus




Atypical





Melanoma in situ





Fibrillar pattern anchored on the ridges (Type B)














Uncommon benign patterns


Globular pattern	Globules not associated with a parallel pattern	4	
Homogeneous pattern	Light brown homogeneous pigmentation with an amorphous appearance	• • •	
Acral reticular pattern	Well-defined pigment network not associated with the skin markings		


Blue nevi on volar surfaces cross ridges & furrows


Often display slight IDP

Monitor















































JUNE 11-13TH | BONUS WORKSHOP ON JUNE 10TH

THE GREENBRIER
NEW RIVER GORGE NATIONAL PARK, WV



REGISTER AT AMERICANDERMOSCOPY.COM



7th World Congress of **DERMOSCOPY 2027**

12-14 May 2027 - Congress Center - Cité Internationale - LYON